arXiv:1408.3227v1 [astro-ph.SR] 14 Aug 2014

A close-in companion to HD142384[1]

J.-B. Le Bouquin
Univ. Grenoble Alpes, IPAG, F-38000 Grenoble, France
CNRS, IPAG, F-38000 Grenoble, France

Prepared 2014/06/25

Abstract: I report the detection of a companion around HD142384 from Optical Long Baseline Interferometry in the near-infrared (flux difference ΔHmag ≈ 4 and separation ≈ 20 mas). HD142384 has been used as PSF reference for published observations of the well-known young star HD142527. This is now strongly discouraged.

Observations

Over the last year, HD142384 has been used as reference star for various observations performed with the PIONIER combiner (Le Bouquin et al. 2011) from the Very Large Telescope Interferometer (VLTI, Haguenauer et al. 2010). By chance, most of these observations were double-calibrated by at least one additional reference star.

Table 1 summarizes the date of observations and the list of additional reference stars that could be used to calibrate the observations of HD142384. These reference stars were found with the `SearchCal`[2] (Bonneau et al. 2011). *A-priori*, these reference star are not more trustable than HD142384 and we are blocked in a chicken-and-egg problem. However two arguments support the idea that HD142384 is the culprit: (1) During good nights, the other reference stars provide consistent transfer function estimates. (2) The transfer function measured on HD142384 is systematically *lower* than those computed with the other reference stars.

Binary fitting

The calibrated observations of HD142384 can be satisfactory adjusted with a binary model. We assume a diameter of 0.5 mas for the primary, according to `SearchCal`, and consider that the secondary is point-like. The best fitting flux ratio is around 2.5%, that is ΔHmag ≈ 4. The best fit binary models for each epoch are shown at the end of the manuscript.

Given the separation and flux ratio, the probability of a contamination by a background /foreground object is extremely small. The orbital motion is detected over one year. This is well compatible with the physical size of the separation (nearly 2AU) according to the Hipparcos parallaxe $\pi = 7.49$ mas.

[1] Based on observations collected with the PIONIER/VLTI instrument at the European Southern Observatory, Paranal, Chile.

[2] http://www.jmmc.fr/searchcal_page

Table 1: Log of observations with the reference stars.

Date	Reference stars
2013-02-19	HD142132
2013-04-21	HD144988
2013-06-03	HD143561, HD144073, HD144475
2013-06-06	HD142132, HD175714
2013-06-14	HD144475, HD142132
2013-06-15	HD142132
2013-06-16	HD142132, HD137598
2013-07-02	HD142132
2014-05-28	HD142132, HD144475
2014-06-02	HD142132, HD144475
2014-06-21	HD142132, HD144475, HD146906

Discussion

HD142384 has been used as reference by Biller et al. (2012) which reports the detection of a low-mass stellar companion to the transitional disk star HD142527 using Spare Aperture Masking (SAM/VLT). HD142384 is the only reference that was observed with the H-band filter. The new companion detected around this reference star (HD142384B, 2.5% in H-band) is significantly brighter than the tentative detection of HD142527b (1.6% in H-band, 0.8% in L-band).

Given its close-in separation, HD142384B has probably insignificant influence on the calibration performed in L-band where the spatial resolution of SAM/VLT is 90 mas. This is confirmed by the trend shown in Fig. 1 from Biller et al. (2012), where all the three reference stars are consistent. In H-band, the situation is not as clear. The spatial resolution of SAM/VLT is 40 mas, that is only twice the possible separation of HD142384B.

Alltogether, the use of HD142384 as reference star is now strongly discouraged, especially with SPHERE or GPI.

Acknowledgments

PIONIER is funded by the Université Joseph Fourier (UJF), the Institut de Planétologie et d'Astrophysique de Grenoble (IPAG), the Agence Nationale pour la Recherche (ANR-06-BLAN-0421 and ANR-10-BLAN-0505), and the Institut National des Science de l'Univers (INSU PNP and PNPS). The integrated optics beam combiner is the result of a collaboration between IPAG and CEA-LETI based on CNES R&T funding. This works made use of the Smithsonian/NASA Astrophysics Data System (ADS) and of the Centre de Donnees astronomiques de Strasbourg (CDS). All calculations and graphics were performed with the freeware `Yorick`.

Bibliography

Biller, B., Lacour, S., Juhász, A., et al. 2012, Astrophysical Journal, Letters, 753, L38

Bonneau, D., Delfosse, X., Mourard, D., et al. 2011, Astron. & Astrophys., 535, A53

Haguenauer, P., Alonso, J., Bourget, P., et al. 2010, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 7734, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series

Le Bouquin, J.-B., Berger, J.-P., Lazareff, B., et al. 2011, Astron. & Astrophys., 535, A67



HD142384 (2013−04−22)



HD142384 (2013−06−04)



HD142384 (2013−06−07)



HD142384 (2013−06−15)




HD142384 (2013−06−15)
vis2
t3Phi
HD142384
mjd=56459.065
min(chi2)=0.5
r=0.03 (0.000)
dE=15.70mas*
dN=−9.78mas*
ρ=18.49mas
θ=121.9deg
ε={5.45mas,3.44mas,103deg}
2013−06−15_SCI_HD142384_oiDataCalib
North (mas) −>
<− East (mas)


HD142384 (2013−06−16)
vis2
t3Phi
HD142384
mjd=56460.084
min(chi2)=1.0
r=0.03 (0.000)
dE=16.02mas*
dN=−9.80mas*
ρ=18.78mas
θ=121.4deg
ε={1.92mas,1.54mas,77deg}
2013−06−16_SCI_HD142384_oiDataCalib
North (mas) −>
<− East (mas)

HD142384 (2013−07−03)



HD142384 (2014−05−29)




HD142384 (2014−06−03)
vis2
t3Phi
HD142384
mjd=56811.102
min(chi2)=1.1
r=0.03 (0.000)
dE=8.12mas*
dN=2.72mas*
ρ=8.56mas
θ=71.5deg
ε={24.74mas,3.63mas,101deg}
2014−06−02_SCI_HD142384_oiDataCalib
North (mas) −>
<− East (mas)


HD142384 (2014−06−22)
vis2
t3Phi
HD142384
mjd=56830.097
min(chi2)=0.9
r=0.03 (0.000)
dE=11.58mas*
dN=0.71mas*
ρ=11.60mas
θ=86.5deg
ε={1.12mas,0.62mas,122deg}
2014−06−21_SCI_HD142384_oiDataCalib
North (mas) −>
<− East (mas)